

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/24235 A2

- (51) International Patent Classification⁷: **A61K 51/04**
- (21) International Application Number: PCT/EP01/05483
- (22) International Filing Date: 11 May 2001 (11.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
00110084.1 12 May 2000 (12.05.2000) EP
- (71) Applicant (for all designated States except US):
MALLINCKRODT, INC. [US/US]; 675 McDonnell
Boulevard, St. Louis, MO 63134 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MAECKE, Helmut, R.** [CH/DE]; Bergfriedweg 7, D-79541 Lorrach (DE).
EISENWIENER, Klaus [DE/DE]; Hauptstrasse 160,
D-79576 Weil (DE). **POWELL, Pia** [CH/CH]; In der
Klosterleben 13, CH-4052 Basel (CH).
- (74) Agents: **VAN SOMEREN, Petronella, Francisca, Hendrika, Maria et al.**; Arnold & Siedsma, Sweelinckplein 1,
NL-2517 GK The Hague (NL).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/24235 A2

(54) Title: PROCHELATORS FOR THE PREPARATION OF RADIOMETAL LABELED MOLECULES HAVING IMPROVED BIOLOGICAL PROPERTIES

(57) Abstract: Chelating compounds for labeling bioactive molecules with a radiometal, having general formula: (I) in which, both Y groups may be positioned either trans as shown or cis; A is an effector molecule, such as a peptide, in particular octreotide, CCK, substance P, gastrin, a protein, in particular an antibody or enzyme, sugars or radiosensitizing agents, like doxorubicin; R is a hydrogen, a C₁-C₃ alkyl or a alcohol; X is a spacer, in particular (CH₂)_n-X', in which n is 1-10 and X' is COOH, NH₂, SH, OH or O-halogen, in which halogen is in particular Br, I or Cl, or a molecule of formula (a) or of formula (b), Y is COO⁻, CH₂CONH₂, CH₂CH₂OH, optionally complexed with a radiometal.

WO 02/24235

PCT/EP01/05483

1

**PROCHELATORS FOR THE PREPARATION OF
RADIOMETAL LABELED MOLECULES HAVING
IMPROVED BIOLOGICAL PROPERTIES**

The present invention relates to a convenient synthesis of novel bifunctional prochelators for coupling to bioactive peptides for radiometal labeling and to the radiometal labeled peptides that can be prepared while
5 using these novel prochelators.

DOTA (1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10 tetraazacyclo dodecane) and its derivatives constitute an important class of chelators for biomedical applications as they accommodate very stably a variety of di- and
10 trivalent metal ions. $Gd(DOTA)^{-1}$ is an important MRI (Magnetic Resonance Imaging) contrast agent and as bifunctional versions DOTA is used in radioimmunotherapy.

An emerging area is the use of chelator conjugated bioactive peptides for labeling with
15 radiometals in different fields of diagnostic and therapeutic nuclear oncology. For their convenient and high yield synthesis, prochelators (compounds which become chelators upon deprotection) are necessary which are compatible with the solid and solution phase peptide
20 synthetic procedures. Up till now such prochelators have not been available.

It is therefore the object of the present invention to provide a novel class of prochelators which are compatible with solid and solution phase peptide
25 synthetic procedures, which prochelators can be used for the coupling of chelators to bioactive effector molecules, such as peptides.

According to the invention bifunctional macrocyclic synthons (prochelators) are provided based on
30 DOTA, TRITA, TETA or structures comprising not 4, but 5 or 6 N atoms, which synthons are differentially protected and compatible with solid phase peptide synthesis procedures for labeling with hard Lewis acid radiometals.


WO 02/24235

PCT/EP01/05483

2

Accordingly, the invention relates to polyazamacrocyclic compounds for radiometal labeling, comprising an N_n system, wherein n is 4, 5 or 6, with varying ring size, and wherein at least one of the N atoms is substituted with a free carboxylate group for coupling to an amino function in a bioactive effector molecule, while all N atoms carry a protected sidechain. After coupling of the bioactive effector molecule the protected sidechains can be deprotected to expose the chelating functions for labeling.

Particular compounds of the invention have the general formula:



20 An example of such a compound is 1-(1-carboxy-3-carbotertbutoxypropyl)-4,7,10(carbotertbutoxymethyl)-1,4,7,10-tetraazacyclododecane (DOTAGA(tBu)4).

Compounds of the invention are prepared as follows. Starting from a relevant amino acid, the α -bromo-derivative thereof is synthesized. This derivative is orthogonally protected (tBu, Bzl). This alkylating agent will be reacted with cyclen, cyclam etc. to form a 1:1 adduct followed by tris-alkylation with bromoacetic acid tert-butylester and catalytic hydrogenation with H_2/Pd .

The synthon is monoreactive, carrying a free carboxylate group for coupling to the N-terminal end of the peptide and can be coupled to any biomolecule which then after deprotection can be labeled with a multitude of radiometals known to the person skilled in the art.


The invention relates more in particular to chelating compounds for radioactive labeling of bioactive

WO 02/24235

PCT/EP01/05483

3


molecules, which chelating compounds have the general formula:



in which:

both Y groups may be positioned either trans as shown or cis;

- 15 A is an effector molecule, such as a peptide, in particular octreotide, CCK, substance P, gastrin, a protein, in particular an antibody or enzyme, sugars or radiosensitizing agents, like doxorubicin;
 R is a hydrogen, a C₁-C₃ alkyl or an alcohol;
 20 X is a spacer, in particular (CH₂)_n-X', in which n is 1-10 and X' is COOH, NH₂, SH, OH or O-halogen, in which halogen is in particular Br, I or Cl
 or X is a molecule of the formula



Y is COO⁻, CH₂CONH₂, CH₂CH₂OH.

- In principle, the invention relates to every
 30 possible combination of the above identified substituents whether or not they are explicitly described herein in a specific combination or not.

- Instead of an N₄ system the compound may comprise a N₅ or N₆ system and can be used for different
 35 natural and unnatural amino acids, such as protected cysteine etc. The list of natural and unnatural amino acids is known to the person skilled in the art.

WO 02/24235

PCT/EP01/05483

4

One of the new DOTA-based bifunctional
prochelators, is 1-(1-carboxy-3-carbotertbutoxypropyl)-
4,7,10(carbotertbutoxymethyl)-1,4,7,10-
tetraazacyclododecane (DOTAGA(tBu)₄ (6d)), which serves
5 herein as a model compound. It should be noted that other
compounds of the invention can be prepared in an
analogous manner. The invention is not limited to
compound 6d alone.

Other embodiments of the invention are listed
10 in Table 3.

The 5 step synthesis of 6d is described in the
Examples. It has an overall yield of about 20%. The
coupling of 6d to a bioactive peptide on solid phase is
exemplified with use of a CCK-B (cholecystokinin)
15 analogue.

Described herein are the synthetic steps
towards bifunctional orthogonally protected prochelators
for coupling to the N-terminus of bioactive peptides or
other useful amino functions in biomedical applications.
20 The DOTA-derived chelator should provide 4 intact
carboxylic acid functions besides the macrocyclic
tetraazacyclododecane ring for a stable and efficient
binding of metal ions and a function for biomolecule
coupling.

The strategy includes the synthesis of an
orthogonally protected bromo-alkyl-dicarboxylic acid
diester for the monoalkylation of cyclen(1,4,7,10-
tetraazacyclododecane). The synthesis of compound 6 (n=1,
2) is a 5 step procedure starting from the commercially
30 available aspartic (1b) or glutamic acid-4-(5) benzyl
ester (1d) (Scheme 1) using a method analogous to
Holmberg (Chemische Berichte 1927, 60, 2197-2205)
followed by tert-butylation using tert-butyltrichloro-
acetimidate (TBTA) as reagent (Armstrong, A. et al.,
35 Tetrahedron Lett. 1988, 29, 2483-2486).

The monoalkylation of cyclen, the crucial step,
showed strongly differing yields depending on the bromo-
alkyl-dicarboxylic acid diester (3a-d) used (Table 1).

WO 02/24235





PCT/EP01/05483

5

Table 1


Mono-alkylation yields of cyclen with different
bromodicarboxylic acid esters

5

	alkyl. agent	yield
10		79%
15		21%
		<5%
		83%

Scheme 1:

Synthesis of α -bromosuccinic acid-1-tertbutylester-4-
20 benzyl ester (3b) and α -bromoglutaric acid-1-tertbutyl
ester-5-benzyl ester (3d).



30

In earlier studies (André et al., Chem. Eur. J. 1999, 5, 2977-2982) the strategy was to use metals as protecting groups. In that work it was attempted to introduce succinic acid-di-tert-butylester (3c) and
35 yields below 5% were found for the monoalkylation with the elimination product fumaric acid-di-tert-butylester as the main product. Interestingly the corresponding diphenylmethyl diester (3a) gave high monoalkylation


WO 02/24235

PCT/EP01/05483

6


yields and negligible elimination. With the homologous 2-bromoglutaric-1-tertbutyl-5-benzylester (3d), no elimination product was found, obviously because no conjugated pi-system could be formed.

5 The remaining nitrogens were alkylated by use of three equivalents of bromoacetic acid-tertbutyl ester in $\text{CHCl}_3/\text{K}_2\text{CO}_3$.



15

Deprotection of the benzyl ester group was performed with $\text{H}_2/\text{Pd/C}$.



The overall yield of 1-(1-carboxy-3-carbotertbutoxypropyl)-4,7,10(carbotertbutoxymethyl)-1,4,7,10-tetraazacyclododecane (DOTAGA(tertBu)₄) (6d) over 5 steps was about 20% and of 1-(1-carboxy-2-carbotertbutoxyethyl)-4,7,10(carbotertbutoxymethyl)-1,4,7,10-tetraazacyclododecane (DOTASA(tertBu)₄) (6b) only about 2%.

The convenient use of 6d is exemplified by its coupling to the CCK-B analogue D-Asp-Tyr-Nle-Gly-Trp-Nle-Asp-Phe-NH₂ (7) attached to Rink-amide resin using HATU (O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluroniumhexafluorophosphate) as coupling reagent. After deprotection, (18h, rt, TFA:phenol:thioanisole:water 85:5:5:5) DOTAGA-7 was obtained in high yield and showed

WO 02/24235

PCT/EP01/05483

7

superior properties in comparison to other radiolabeled CCK-B analogues.

According to the invention it was thus found that the new type of prochelators in general and prochelator **6d** in particular, have widespread utility in the field of metalloradiopeptides, other radiolabeled biomolecules and for the synthesis of Gd^{3+} based MRI contrast agents. DOTAGA will allow to label with different radiometals for both diagnostic (^{111}In , $^{67/68}Ga$) and internal radiotherapeutic applications (^{90}Y , ^{177}Lu).

The present invention will be further illustrated in the examples that follow. Synthetic routes for the preparation of the compounds of the invention are given in Figures 1-3.

15

EXAMPLES

EXAMPLE 1

Typical procedure for the reaction of 1 to 2

To a solution of 6 g (25.9 mmol) L-glutamic acid-5benzylester (**1d**) and 9.1 g (88.5 mmol) sodium bromide in 45 ml aqueous 1N hydrobromic acid (46 mmol) cooled to 0°C was added portionwise 3.175 g (46 mmol) sodium nitrite. After stirring for 2h at 0°C 2.25 ml concentrated sulfuric acid was added followed by diethylether.

The water phase was extracted 3 times with diethylether. The combined organic phases were extracted 4 times with brine, dried over Na_2SO_4 and concentrated.

The crude product was purified by chromatography (silica gel 60; hexane/EtOAc 3:1 to 2:1) and obtained as a yellow oil in a yield of 4.8 g (63%).
 1H -NMR (300 MHz, $CDCl_3$, $SiMe_4$): 10.1 (1H, $COOH$); 7.3 (m, 5H, Ar); 5.15 (s, 2 H, CH_2 -Ph); 4.4 (dd, $^3J = 5.7$, 1H, $CHBr$); 2.6 (t, $^3J = 6.8$, 2H, CH_2 - $COOBzl$); 2.5-2.2 (m, 2H, $CHBr$ - CH_2 - CH_2); ^{13}C -NMR (75 MHz, $CDCl_3$, $SiMe_4$): 174.5 ($COOH$); 171.9 ($COOBzl$); 135.5 ($CH_2C(Ar)$); 128.6, 128.4, 128.3 ($C(Ar)$); 66.8 ($O-CH_2-Ar$); 44.1 ($HCHBr$); 31.4 ($HCHBr-CH_2$);

WO 02/24235

PCT/EP01/05483

8

29.4 (CH_2COOBzl ; EI-MS m/z (intensity): 302, 300 (12, $[\text{M}]^+$); 91 (100, $[\text{Bzl}]^+$).

5 EXAMPLE 2

Typical procedure for the reaction of 2 to 3

To a solution of 4.8 g (15.9 mmol) 2d in 20 ml CHCl_3 a solution of 6.26 ml (34.1 mmol) TBTA (tert-butyltrichloroacetimidate) in 20 ml cyclohexane was added dropwise over 20 min. During the addition a white precipitate formed, which was dissolved by the addition of 3.5 ml of DMA followed by 320 μl boron trifluoride ethyl etherate as catalyst. The reaction mixture was stirred for 3 days at Room Temperature (RT).

The mixture was concentrated and the remaining DMA phase was extracted 3 times with 30 ml hexane. The hexane phase was evaporated and the residue chromatographed over silica gel 60 (Hexane/EtOAc 20:1 later 9:1) affording 3.5 g (61%) of a colourless liquid.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , SiMe_4): 7.4 (m, 5H, Ar); 5.15 (s, 2H, $\text{CH}_2\text{-Ph}$); 4.35 (dd, 1H, CHBr); 2.6 (td, 2H, $\text{CH}_2\text{-COOBzl}$); 2.5-2.2 (m, 2H, $\text{CHBr-CH}_2\text{-CH}_2$); 1.5 (s, 9H, $\text{C}(\text{CH}_3)_3$). $^{13}\text{C-NMR}$ (75 MHz, CDCl_3 , SiMe_4): 172.4 (COOBzl); 168.7 (COOtBu); 136.1 ($\text{CH}_2\text{C}(\text{Ar})$); 129.0, 128.8, 128.7 ($\text{C}(\text{Ar})$); 83.1 ($\text{C}(\text{CH}_3)_3$); 67.0 ($\text{OCH}_2\text{-Ar}$); 47.1 (HCBrl); 32.0 (HCBrl-CH_2); 30.1 (CH_2COOBzl); 28.1 ($\text{C}(\text{CH}_3)_3$); EI-MS m/z (intensity): 302, 300 (18, $[\text{M-C}_4\text{H}_9]^+$); 57 (100, $[\text{C}_4\text{H}_9]^+$).

30 EXAMPLE 3

General procedure of the monoalkylation of cyclen

A solution of 870 mg (2.44 mmol) a-bromoglutaric acid-1-tert-butylester-5-benzylester (3d) in CHCl_3 was added dropwise over a period of 1h to a solution of 885 mg (4.9 mmol) cyclen in 4 ml CHCl_3 . The mixture was stirred for 2 days at room temperature and concentrated to a brown oil. The crude product was purified by column chromatography (silica gel 60;

WO 02/24235

PCT/EP01/05483

9

ethanol/NH₃ 95:5), yield 920 mg (83%) of a colourless oil.

¹H-NMR (300 MHz, CDCl₃, SiMe₄): 7.35 (m, 5H, Ar); 5.1 (s, 2H, CH₂-Ph); 3.25 (dd, 1H, CHBr); 2.9-2.5 (m, 18H, NCH₂, CH₂COOBzl); 2.2-1.85 (m, 2H, CHN-CH₂-CH₂); 1.45 (s, 9H, C(CH₃)₃); ¹³C-NMR (75 MHz, CDCl₃, SiMe₄): 173.1 (COOBzl); 171.5 (COOtBu); 135.8 (CH₂C(Ar)); 128.5, 128.3, 128.2 (C(Ar)); 81.4 (C(CH₃)₃); 66.2 (O-CH₂-Ar); 63.5 (HCNCH₂); 48.8, 48.0, 46.5, 45.6 (NCH₂CH₂N); 30.6 (CH₂COOBzl); 28.2 (C(CH₃)₃); 24.5 (HCN-CH₂); EI-MS m/z: (intensity): 449.3 (56, [M+H]⁺); 245.8 (100, [M+CH₃CN+2H]⁺⁺).

EXAMPLE 4

15 Synthesis of 1-(1-carbobenzyloxy-3-carbotertbutoxy-propyl)-4,7,10-(carbotertbutoxymethyl)-1,4,7,10-tetraazacyclododecane (5d)

A suspension of 1.1 g (5.6 mmol) bromoacetic acid-tert-butylester, 1.02 g (2.27 mmol) 1-(1-carbo-
benzyloxy-3-carbotertbutoxypropyl)1,4,7,10-tetraaza-
cyclododecane (4d), and 2.63 g (19.1 mmol) of dry
potassium carbonate in 10 ml dry acetonitrile was stirred
for 18 h at RT and was filtrated afterwards over Celite
and evaporated to dryness.

The crude product was purified by column
chromatography (silica gel 60; CH₂Cl₂/EtOH 9:1 followed by
EtOH/NH₃ 95:5) yield 1.3 g (73%) of a yellow oil (5d). ¹H-
NMR (300 MHz, CDCl₃, SiMe₄): 7.35 (m, 5H, Ar); 5.1 (s, 2H,
CH₂-Ph); 3.6-1.9 (m, 27H, CHN, NCH₂, CH₂COOBzl, CHNCH₂-CH₂,
CH₂COOC(CH₃)₃); 1.45 (s, 36H, C(CH₃)₃); ¹³C-NMR (75 MHz,
CDCl₃, SiMe₄): 174.6 (COOBzl); 172.9, 172.8, 172.6
(COOtBu); 135.6 (CH₂C(Ar)); 128.5, 128.3, 128.2 (C(Ar));
82.4, 81.8, 81.8 (C(CH₃)₃); 66.3 (O-CH₂-Ar); 55.8, 55.7,
55.4, 52.6, 52.3, 50.3, 48.5, 48.1, 47.1, 44.3 (13C,
HCNCH₂, NCH₂CH₂N, CH₂COOtBu, CH₂COOBzl); (NCHCH₂CH₂); 28.0,
28.0, 27.8, 27.6 (C(CH₃)₃); EI-MS m/z (intensity): 813.6
(22, [M+Na]⁺); 791.6 (38, [M+H]⁺); 396.5 (100, [M+2H]⁺⁺).

WO 02/24235

PCT/EP01/05483

10

EXAMPLE 5Synthesis of DOTAGA(tBu)₄ (6d)

600 mg (0.76 mmol) 5d was dissolved in methanol, and 30 mg Pd/C suspended in 1 ml H₂O was added. 5 The mixture was hydrogenated for 2 days, filtrated over Celite and evaporated to dryness. The crude product was chromatographed on silica gel 60 (EtOH/NH₃ 95:5) to obtain 470 mg (84.6%) of a white solid (6d). ¹H-NMR (300 MHz, CDCl₃, SiMe₄): 6.5 (br, 1H, COOH); 3.6-2.0 (m, 27H, CHN, NCH₂, CH₂COOH, CHN-CH₂-CH₂, CH₂COOC(CH₃)₃); 1.45 (s, 36H, C(CH₃)₃); ¹³C-NMR (75 MHz, CDCl₃, SiMe₄): 175.2 (COOH); 175.0, 172.9, 172.8, 172.6 (COOtBu); 82.4, 82.1, 81.9 (C(CH₃)₃); 55.8, 60.1 (NCHCOOtBu); 55.9, 55.8, 55.6, 52.7, 52.6, 52.5, 48.6, 48.5, 48.2, 47.1, 44.3 (12C, NCH₂CH₂N, CH₂COOtBu, CH₂COOH); 33.4 (NCHCH₂CH₂); 27.9, 27.8 (C(CH₃)₃); EI-MS m/z (intensity): 723.5 (27, [M+Na]⁺); 701.5 (68, [M+H]⁺); 351.4 (100, [M+2H]⁺⁺).

20 EXAMPLE 6Preparation of DOTAGA-7

Compound 6d was coupled to the CCK-B analogue D-Asp-Tyr-Nle-Gly-Trp-Nle-Asp-Phe-NH₂ (7) attached to Rink-amide resin using HATU (O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluroniumhexafluorophosphate) as coupling reagent. 25

After deprotection, (18h, rt, TFA:phenol:thioanisol:water 85:5:5:5) DOTAGA-7 was obtained in high yield and showed superior properties in comparison to other radiolabeled CCK-B analogues. 30

The data of the DOTAGA-CCK-B analogue (DOTAGA-7) were as follows: Yield: 12.7 mg, HPLC purity >95%, (+) EI-MS m/z (intensity): 1486.1 (48, [M+H]⁺); 743.7 (60, [M+2H]⁺⁺); (-) EI-MS m/z (intensity): 1484.0 (28, [M+H]⁻); 35 741.8 (90, [M+2H]⁻⁻).

WO 02/24235

PCT/EP01/05483

11

EXAMPLE 7**Biological properties**

It has been found that when using the
prochelators of the invention for preparing biologically
5 active molecules, these molecules have better biological
properties than molecules prepared with other
(pro)chelators. The advantages for example are a better
labeling yield according to the following table 2:

10 Table 2

	amount of radioactive label	labeling efficiency of DOTATOC- peptide	labeling efficiency of DOTA3-peptide (DOTAGA) (invention)
5 µg of the compound to be labeled	5 mCi ⁹⁰ Y	99.5%	99.9%
	10 mCi ⁹⁰ Y	95%	100%
	20 mCi ⁹⁰ Y	92%	99.9%

15

In addition, the stability is higher. The reaction of ⁹⁰Y-
DOTATOC or ⁹⁰Y-DOTAGA at 37°C with the chelator DTPA
results in ⁹⁰Y-DTPA. The half life of this reaction is 23
hours for DOTATOC and 79 hours for DOTAGA.

20

Table 3 shows various biological properties of
the compound of the invention Y-DOTA3TOC and Y-
DOTA13TOC. The radiometal is not shown.

WO 02/24235

PCT/EP01/05483

12

Biological results of Yttrium labelled Peptides







Peptide	IC ₅₀ (hsst2)	Tumor	Kidney	Charge
 <p>Y-DOTATOC: Y-DOTATyr³octreotide</p>	11 ± 1.7	13.5	12.3	+1
adrenal, pancreas low				
 <p>Y-DOTATATE: Y-DOTATyr³octreotate</p>	1.6 ± 0.4	14.5	8	0
 <p>Y-DOTA3TOC: Y-DOTA3tyr³octreotide</p>	1.5 ± 0.5	30	56	0
 <p>Y-DOTA3TATE: Y-DOTA3tyr³octreotate</p>	3.5	13.5	68	-1
adrenal, pancreas high				
 <p>Y-DOTA13TOC: Y-DOTA13tyr³octreotide</p>	28 ± 10	23.5	25.5	0
 <p>Y-DOTA13TATE: Y-DOTA13tyr³octreotide</p>	0.23			+1

Table 3

WO 02/24235


PCT/EP01/05483

13

CLAIMS


1. Polyazamacrocyclic compounds for radiometal labeling, comprising an N_n system, wherein n is 4, 5 or 6, with varying ring size, and wherein at least one of the N atoms is substituted with a free carboxylate group for coupling to an amino function in a bioactive effector molecule, while all N atoms carry a protected sidechain.

2. Compound as claimed in claim 1 having the
10 general formula:



3. Compound as claimed in claim 1 or 2, which
compound is 1-(1-carboxy-3-carbotertbutoxypropyl)-
4,7,10(carbotertbutoxymethyl)-1,4,7,10-
25 tetraazacyclododecane (DOTAGA(tBu)4).

4. Chelating compounds for labeling bioactive
molecules with a radiometal, having the general formula:



in which:

both Y groups may be positioned either trans as shown or
cis;

WO 02/24235


PCT/EP01/05483

14


A is an effector molecule, such as a peptide, in particular octreotide, CCK, substance P, gastrin, a protein, in particular an antibody or enzyme, sugars or radiosensitizing agents, like doxorubicin;

5 R is a hydrogen, a C₁-C₃ alkyl or a alcohol;

X is a spacer, in particular (CH₂)_n-X', in which n is 1-10 and X' is COOH, NH₂, SH, OH or O-halogen, in which halogen is in particular Br, I or Cl or a molecule of the formula



or of the formula



Y is COO⁻, CH₂CONH₂, CH₂CH₂OH, optionally complexed with a radiometal.

20 5. Compounds as claimed in claim 4, wherein R is hydrogen, n is 1, X' is COOH, Y is COO⁻ and A is as defined in claim 3.

6. Compound as claimed in claim 5, wherein R is hydrogen, n is 1, X' is COOH, Y is COO⁻ and A is
25 octreotide or octreotate.

7. Compound as claimed in claim 4, wherein R is COOH, n is 1, X' is COOH, Y is COO⁻ and A is as defined in claim 3.

8. Compound as claimed in claim 7, wherein R is
30 COOH, n is 1, X' is COOH, Y is COO⁻ and A is octreotide or octreotate.

9. Compounds as claimed in claim 4, selected from the group consisting of DOTATyr³octreotide, DOTATyr³octreotate, DOTA3tyr³octreotide,
35 DOTA3tyr³octreotate, DOTAT3tyr³octreotide, DOTATa.13tyr³octreotate.

WO 02/24235

PCT/EP01/05483

15

10. Use of compounds as claimed in claims 1-3
for the preparation of compounds as claimed in claims 4-
9.

11. Method for the preparation of radiometal
5 labeled bioactive molecules, comprising the steps of:

a) synthesizing compounds as claimed in claims
1-3 having protected side chains on the N atoms and a
free carboxylate group;

b) coupling a bioactive molecule to the free
10 carboxylate group;

c) deprotecting the protected side chains; and

d) labeling the chelator structure thus
obtained with a desired radiometal.


12. Compounds as claimed in claims 4-9 labeled
15 with a radiometal for use in diagnosis and therapy.

13. Use of compounds as claimed in claims 4-9
labeled with a radiometal for the preparation of a
diagnostic or therapeutical composition for treatment of
various diseases.

20 14. Use as claimed in claim 13, wherein the
radiometal label is ^{90}Y .


WO 02/24235

PCT/EP01/05483

**FIG. 1A**


WO 02/24235

PCT/EP01/05483



WO 02/24235


PCT/EP01/05483



WO 02/24235

PCT/EP01/05483

4/5

FIG. 2B

WO 02/24235

PCT/EP01/05483

5/5




FIG. 3